Medicinal Ingredients (per serving)

Vitamin C (Sodium ascorbate)	500mg
Phosphatidylcholine (Glycine max-seed)	157mg
R-alpha-Lipoic acid	25mg

Non-Medicinal Ingredients: Water, glycerin, ethanol, lemon peel essential oil

Recommended Use: Helps to maintain immune function.

Recommended Dose: Adults: Take 4 pumps (2 mL) daily, holding for 30 sec. before swallowing. May also be stirred into water. Best taken on an empty stomach at least 10 minutes before meals. Use within 30 days of opening.

Cautions and Warnings: Consult a health care practitioner prior to use if you are pregnant, breastfeeding or have diabetes.

Known Adverse Reactions: If you experience symptoms of hypoglycemia (serious low blood sugar) including paleness, chills, confusion, feelings of anxiety, dizziness, tremor, sweating, nausea or headache, discontinue use and consult a health care practitioner.

Refrigerate upon receipt.

Voir à l'intérieur pour le français

Manufactured by/Fabriqué par: Quicksilver Scientific
1960 Cherry St.
Louisville, CO 80027

Importer/Importateur
DeVita Vite Ltd.
Box 283 Keswick ON
L4P 3E2 (800) 840-1676

QUICKSILVER DELIVERY SYSTEMS®

Product Code
Code produit: QSCRLA02
15067.04

Lot:

Exp.:

7 93888 08211 3

QUICKSILVER SCIENTIFIC®

LIPOSOMAL

VITAMIN C + R-LIPOIC ACID

VITAMIN C LIPOSOMALE CONTIENT DE L'ACIDE R-LIPOÏQUE

NPN 80079588

Natural Health Product
Produit de santé naturel

50 ML LIQUID/(LIQUIDE)

PEEL HERE ▶

HINGE

Ingrédients médicinaux (par portion)

Vitamine C (ascorbate de sodium) 500mg
Phosphatidylcholine (graine de Glycine max) 157mg
Acide R-alpha-lipoïque 25mg

Ingrédients non médicinaux : Eau, glycérine, éthanol, huile essentielle d'écorce de citron

Emploi recommandé : Aide à maintenir le fonctionnement du système immunitaire.

Dose recommandée : Adultes : 4 pompes (2ml) par jour. Garder dans la bouche pendant 30 secondes avant d'avaler. Peut aussi être mélangé à de l'eau. Il est recommandé de prendre le produit à jeun 10 minutes au moins avant les repas. A utiliser dans les 30 jours après ouverture.

Avertissements et mises en garde : Consultez un professionnel de la santé avant d'utiliser ce produit si vous être enceinte, si vous allaitez ou si vous avez le diabète.

Réactions indésirables connues : Si vous présentez des symptômes d'hypoglycémie (glycémie très basse), y compris de la pâleur, des frissons, de la confusion, de l'anxiété, des étourdissements, des tremblements, des sueurs, des nausées ou des maux de tête, cessez d'utiliser le produit et consultez un professionnel de la santé.

Mettre au réfrigérateur dès la réception.